

Research Article OPEN ACCESS

Lisnerová M., Bartošová-Sojková P., Lövy A., Blabolil P., Fiala I. 2024: Evidence of striking morphological similarity and tissue tropism of phylogenetically distant myxozoan genera: *Myxidium* and *Paramyxidium* in the kidney of the European eel. Folia Parasitol. 71: 013.

Supplementary Table 1. List of primer pairs used for PCR amplification of sequences of *Myxidium giardi* Cépède, 1906 and for screening fish for *M. giardi* infections.

Primer name	Sequence (5'→3')	Annealing temperature (°C)	Length of amplified PCR product	References
18e	TGGTTGATCCTGCCAGT	64	~ 2,000 bp	Hillis and Dixon (1991)
18g	GGTAGTAGCGACGGGCGGTGTG			
Myxgp2f	WTGGATAACCGTGGGAAA	58	~ 900 bp	Kent et al. (1998) Hallett and Diamant (2001)
ACT1r	AATTTACCTCTCGCTGCCA			
MgiarF	GAACAGTTGTGTTTATGTGC	60	~1,100 bp	Present study Hillis and Dixon (1991)
18g	GGTAGTAGCGACGGGCGGTGTG			

Address for correspondence: Martina Lisnerová, Institute of Parasitology, Biology Centre of the Czech Academy of Sciences, Branišovská 31, 370 05 České Budějovice, Czech Republic. E-mail: lisnerova@paru.cas.cz; ORCID-iD 0000-0003-3565-4816